

Occurrence of blackleg and current preventive measures against outbreaks in the Canton of Bern

L. Käser¹, R. Huber³, S. Häsler⁵, S. Kittl⁴, B. Vidondo², H. Posthaus¹, I. Cvitas¹

¹Institute of Animal Pathology, Vetsuisse-Faculty, University of Bern, Switzerland; ²Veterinary Public Health Institute, Vetsuisse-Faculty, University of Bern, Switzerland; ³Cantonal Veterinary Office, Bern, Switzerland; ⁴Institute of Veterinary Bacteriology, Vetsuisse-Faculty, University of Bern, Switzerland; ⁵Swiss Association for Veterinary History, Bern, Switzerland

<https://doi.org/10.17236/sat00342>

Eingereicht: 26.07.2024
Angenommen: 24.10.2024

Auftreten von Rauschbrand und aktuelle Präventionsmassnahmen gegen Ausbrüche im Kanton Bern

Rauschbrand ist eine akute, tödlich verlaufende Infektionskrankheit, die durch *Clostridium chauvoei* verursacht wird. Sie betrifft hauptsächlich Rinder und ist durch schwere nekrohämorrhagische Myositis und Myokarditis gekennzeichnet. Während die Krankheit in der Vergangenheit erhebliche wirtschaftliche Verluste verursachte, ist die Zahl der erkrankten Rinder in den letzten 30 Jahren zurückgegangen. Daraufhin wurde der Seuchenstatus von Rauschbrand in der Schweiz offiziell geändert und die Krankheit ist seit 2022 nicht mehr meldepflichtig. Angesichts der möglichen Persistenz von *C. chauvoei*-Sporen in der Umwelt kann Rauschbrand in bestimmten Gebieten des Kantons Bern, einer wichtigen und grossen Agrarregion der Schweiz, jedoch weiterhin auftreten.

In dieser Studie haben wir die epidemiologischen Daten zu Rauschbrand von 1992–2023 und die aktuell durchgeführten prophylaktischen Massnahmen der Tierärzte in den Risikogebieten des Kantons Bern ausgewertet. Unsere Ergebnisse zeigen, dass die meisten Fälle in Gebieten auftraten, die vom kantonalen Veterinäramt weiterhin als Risikogebiete eingestuft wurden. Jedoch wurden auch vereinzelt Rauschbranderkrankungen in Gebieten, die seit 2020 nicht mehr als Risikogebiete eingestuft sind, diagnostiziert. Gleichzeitig meldeten Tierärzte in den letzten Jahren Verdachtsfälle sowie eine verringerte Akzeptanz der Impfung bei Landwirten in Risikogebieten.

Die Kombination aus fehlender Überwachung, vermindertem Seuchenbewusstsein, sinkender Impfbereitschaft der Landwirte und der langen Umweltpersistenz von *C. chauvoei*-Sporen stellt ein Risiko für ein erneutes Auftreten von Rauschbrand dar. Tierärzte, die in ehemals betroffenen Gebieten im Kanton Bern und anderswo in der Schweiz, praktizieren, sollten bei einem Verdacht auf Rauschbrand entsprechend handeln.

Schlüsselwörter: Rauschbrand, Kanton Bern, Rind, *Clostridium chauvoei*, Risikogebiet, Impfakzeptanz

Abstract

Blackleg is an acute, lethal, infectious disease caused by *Clostridium chauvoei*. It affects mainly cattle and is characterized by severe necrohemorrhagic myositis and myocarditis. While the disease has caused substantial economic losses in the past, the number of diseased cattle has been decreasing in the last 30 years. Subsequently, the status of blackleg has been officially changed in Switzerland and the disease is no longer reportable since 2022. Considering the potential persistence of *C. chauvoei* spores in the environment, blackleg may however still be prevalent in certain areas of the Canton of Bern, an important and large agricultural region of Switzerland. In this study, we evaluated both the spatio-temporal distribution of blackleg from 1992–2023 and the currently implemented prophylactic measurements used by veterinarians in the risk areas of the Canton of Bern. Our results show that most of the cases occurred in areas still defined as risk areas by the Cantonal Veterinary Office, however some occurred in areas no longer listed as risk areas since 2020. In parallel, veterinarians reported suspected cases in recent years as well as a decreased acceptance of vaccination by farmers in risk areas.

The combination of lack of blackleg monitoring, reduced disease awareness, decreasing willingness of farmers to vaccinate and the long environmental persistence of *C. chauvoei* spores represent a risk for reoccurrence of the disease. Veterinarians practicing in once affected areas, in the Canton of Bern and elsewhere in Switzerland, should be aware of this and act accordingly once blackleg outbreaks are suspected.

Keywords: Blackleg, Canton of Bern, cattle, *Clostridium chauvoei*, risk area, vaccine acceptance

Introduction

Clostridium chauvoei, a gram-positive, anaerobic, rod-shaped, spore-forming bacterium, is the causative agent of blackleg, an acute, infectious disease causing significant losses in cattle but also other ruminants and wild animals worldwide.^{8,16} Cattle aged between six and 24 months are most commonly affected.^{1,9} Clinically, blackleg is characterized by a severe necrotizing and hemorrhagic myositis,¹ affecting mostly muscles of the hindlimbs, chest, diaphragm, heart and tongue,⁷ fever and ultimately rapid death of the affected animal. In the peracute form, affected animals are most commonly found dead without any previous signs of illness.⁷ In the acute form, swelling and crepitus of the affected muscles are typical clinical signs.¹⁷ Animals usually get infected with *C. chauvoei* through oral uptake of spores from contaminated soil, water, feed, or less frequently, through wound infection.¹⁷ The spores have the ability to cross the intestinal mucous membrane, and subsequently distribute via the bloodstream to several tissues, including skeletal and cardiac muscle.² There they can remain dormant until a localized hypoxic or anoxic event, e.g. through local trauma, occurs. The establishment of anaerobic conditions enables germination of the spores, subsequent bacterial proliferation, and production of toxins, ultimately leading to muscular necrosis and hemorrhage.⁷ Environmental contamination is mainly caused by spores released from carcasses of animals that succumbed to blackleg.¹⁷ These spores are highly resistant to environmental conditions and can persist in the environment for extended periods of several years to potentially decades.¹

In Switzerland, blackleg occurs endemically. During the 19th and 20th century, it was the cause of substantial economic losses.¹⁴ Most cases in Switzerland were reported in the following regions: Mountains of Bernese Oberland, Alps of the cantons of Fribourg, Jura, Schwyz, Lucerne, St. Gallen, Vaud, and Valais.¹² Until 2009, the disease had a status of “epizootic disease to be controlled”, as defined in the animal disease ordinance (Animal disease ordinance, Art. 4 and 5). This implied for the Canton of Bern that a vaccination program for animals in defined risk areas was financed by the animal disease fund, which is kept by the canton. Cattle between two months and three years of age were vaccinated between February and June. Furthermore, suspected cases were reported to the Cantonal Veterinary Office for surveillance and bacteriologically or epidemiologically confirmed cases were reported publicly every other week.

Risk areas have been defined since the early 20th century. A list of risk areas has been created and continuously updated by the Cantonal Veterinary Office based on locations where cases occurred. Due to decreasing number of cases, the status of the disease in Switzerland has been changed to “epizootic disease to be monitored” in 2009. Thus, in the Can-

ton of Bern, vaccination and testing were not financially supported anymore, however, these were still highly recommended for cattle that were transferred to known risk areas. In addition, animal losses were no longer compensated by the animal disease fund. Confirmed cases were still reported publicly every week in the animal disease notification system (InfoSM), the data bank of the Federal Food Safety and Veterinary Office (FSVO).¹³

Since November 2022, blackleg has not been listed as an epizootic disease in Switzerland anymore. Therefore, cases occurring after November 2022 are not reported and control of blackleg is in the responsibility of the livestock owners.¹⁶ Risk areas, as defined by the Cantonal Veterinary Office, are still publicly listed,¹¹ but have not been updated since 2020. Vaccination is recommended in areas where blackleg has occurred in the past, however, all costs must be covered by the owners. Currently, a multivalent clostridial vaccine, Bovilis Bravoxin* (MSD Animal Health GmbH, Lucerne), is licensed for Switzerland (TAM, swissmedic.ch). This vaccine contains toxoid antigens of *Clostridium perfringens* type A, B, C, D, *Clostridium novyi*, *Clostridium septicum*, *Clostridium tetani*, *Clostridium sordellii* and *Clostridium haemolyticum*, as well as a formaldehyd and thiomersal inactivated whole cell bacterial culture of *Clostridium chauvoei*.¹⁰

Considering the potential persistence of *C. chauvoei* spores in the environment, we hypothesized that blackleg is still prevalent in certain areas of the Canton of Bern, an important and large agricultural region of Switzerland.⁶ Therefore, we evaluated the epidemiological situation of blackleg in the Canton of Bern from 1992 to 2023. In addition, we performed a survey to determine current prophylactic practices used by veterinarians in blackleg risk areas in the Canton of Bern.

Material and Methods

Study area

This study focuses on the Canton of Bern, Switzerland, with a surface area of 5'960 km², including approximately 1'500 Alpine pastures.⁶ The canton is divided into four court regions: Bernese Jura Seeland, Emmental Oberaargau, Bern-Mittelland, and Bernese Oberland.⁴ Within the Canton of Bern, cattle are often moved between different areas. Emmental Oberaargau counts approximately 53'000 cattle, Oberland approximately 49'000, Mittelland 44'000 and Bernese Jura Seeland around 26'000 cattle.⁶

Data collection

The data on officially reported blackleg cases in the Canton of Bern has been collected from the data bank of the Federal Food Safety and Veterinary Office (FSVO), the InfoSM reporting system,¹³ and the Institute of Veterinary Bacteriology (IVB, Vetsuisse-Faculty, University of Bern, Switzer-

Occurrence of blackleg and current preventive measures against outbreaks in the Canton of Bern

L. Käser et al.

Occurrence of blackleg and current preventive measures against outbreaks in the Canton of Bern

L. Käser et al.

land). The blackleg cases were stratified by location with geographical coordinate and date of the case. Risk areas were defined by the Cantonal Veterinary Office based on the occurrence of cases in the past. A list of these areas was updated until 2020. Based on the geographical coordinates, risk areas and confirmed blackleg cases were transferred to geographical maps.

Maps

The creation of the maps was conducted using an open-source geographical information system (QGIS 3.30, 's-Hertogenbosch. qgis.org). To simplify the case occurrence on the maps, the years between 1992 and 2023 were stratified into the following four time periods: 1992–1999, 2000–2009, 2010–2019, 2020–2023. Seven cases could not be precisely located due to lack of information about the exact location of the animal investigated and are therefore not shown on the map.

Questionnaire

Veterinarians practicing in the Canton of Bern were asked to complete a questionnaire (available on request from au-

thors). In total 18 veterinarians, 15 of them actively practicing, three recently retired, took part in the study. Fifteen of them were practicing in the risk areas, whereas three have been practicing in areas bringing animals to alpine pastures in risk areas. The selection of the veterinarians for the study was conducted in collaboration with the Cantonal Veterinary Office.

The questionnaire was split in four parts: the first part involved questions regarding the veterinary practice participants are employed at, the second part comprised questions about clinical blackleg cases the practicing veterinarians have seen, and the third and fourth part involved questions regarding the preventive measures, including current vaccination plans.

It contained 16 closed and six open-ended questions and on average, it took participants approximately 10 minutes to complete. A paper version of the questionnaire was produced, which was used to interview the participants. Out of 18 interviews, 17 were held in the presence of the participating person and the first author, one was conducted via telephone. The interviews were performed between November 2023 and May 2024.




Figure 1: Number of officially reported blackleg cases per year in the Canton of Bern from 1992–2023. Color code corresponds to defined time periods: 1992–1999; 2000–2009; 2010–2019; 2020–2023.

Data analysis

Descriptive data analysis of the cases as well as encoding of the questionnaire was conducted in Microsoft Excel 2024. The responses of the questionnaire were entered manually. NCSS 12 Software (2024, NCSS, LLC. Kaysville, Utah, USA, ncss.com/software/ncss) was used to perform descriptive statistics. For the analysis of the questionnaire, frequency tables were done.

Results

Temporal and spatial distribution of reported blackleg cases in the Canton of Bern from 1992–2023

Altogether, 45 cases of blackleg were reported during the period of 1992–2023. Most cases occurred between 2010 and 2019, however, five cases were reported from 2020 to 2022. In nine out of 30 years, no cases were officially reported (Figure 1).

In the year of 2023, no cases of blackleg were detected in the laboratory of IVB in Bern. Furthermore, due to changes in the epidemiological status of the disease, FSVO has no information on the possible occurrence of blackleg in 2023.

Most cases were reported during summer, from June to September, which coincides with the Alpine pasture period (Figure 2).

Plotting each reported case with clear localization data of the succumbed animal onto maps with published risk areas shows that most of the cases ($n = 30$) occurred in those areas still listed as risk areas in 2023 (Figure 3). However, since 2020, three cases occurred in areas that were no longer listed as risk areas in 2023. These areas were still listed in 2019, but were subsequently removed.

Occurrence of blackleg and current preventive measures against outbreaks in the Canton of Bern

L. Käser et al.




Figure 2: Monthly occurrence of officially reported blackleg cases in the Canton of Bern from 1992–2023.




Figure 3: Spatial distribution of officially reported blackleg cases in the Canton of Bern from 1992–2023. Currently published risk areas (2024, last update 2020) are marked in orange. The line pattern marks the risk areas published until 2020, which were subsequently removed from the list. The stars mark blackleg cases occurring between 1992–2022 in the Canton of Bern color-coded into defined time periods in which cases have occurred (1992–1999; 2000–2009; 2010–2019; 2020–2023).

Occurrence of blackleg and current preventive measures against outbreaks in the Canton of Bern

L. Käser et al.

Results of the survey

Respondent demographics

In total, 18 veterinarians working in 18 different veterinary practices took part in the survey. Participants were actively practicing (n=15), or recently retired veterinarians (n=3). Fifteen out of 18 veterinarians practiced in at least one of the defined risk areas. Three did not have any risk areas within their practice area (Table 1), however, they performed vaccinations in non-risk areas for animals that were seasonally transferred to alpine pastures in declared risk areas. The average age of the participating veterinarians was 58 years, with the oldest being 80 years and the youngest 34 years old.

Occurrence of cases

Fourteen veterinarians (73,68%) reported to have seen at least one clinical case of blackleg in their career, that include bacteriologically confirmed as well as clinically suspected cases. Eight of them only saw cases sporadically (sporadic was defined by the author as less than five per career so far, Table 2), six have seen frequent cases up to several per year over a period of time. Overall, the occurrence of suspected blackleg cases has been reported in all time periods from 1992–2023, with the highest proportion of 40,74% between 2010–2019 (participants were able to select more than one answer). Importantly, four veterinarians (12,90%) saw suspected cases in 2023, which, due to change in epidemiological status of the disease, were not followed up. Frequent cases were only described in the Oberland region.

Table 1: Results of a questionnaire to 18 veterinarians practicing in Canton of Bern in relation to size of the practice area, suspected blackleg cases and case frequency as well as measures against blackleg. Results are displayed in absolute and relative numbers.

Number and % of veterinarians practicing per officially defined risk areas		
	Count	Percent
Risk area	15	83,33%
No risk area	3	16,67%
Number and % of veterinarians and the size of farms (as number of cattle per farm in four categories)		
0–100	4	23,53%
101–300	7	41,18%
301–500	4	23,53%
>500	2	11,76%
Number and % of veterinarians according to the percentage of cattle from farms going to summer pastures in four categories		
0–25%	2	11,76%
26–50%	2	11,76%
51–75%	3	17,65%
76–100%	10	58,82%
Number and % of veterinarians according to whether they observed blackleg case or not		
Observed blackleg case	14	73,68%
Did not observe blackleg case	4	21,05%
Number and % of suspected blackleg cases reported by 18 veterinarians working in Canton of Bern in different time periods		
1992–1999	7	22,58%
2000–2009	3	9,68%
2010–2019	11	35,48%
2020–2022	6	19,35%
2023	4	12,90%
Case frequency (n)		
Regular	6	42,86%
Sporadic (< 5 cases per career)	8	57,14%
Number and % of veterinarians reporting different types of measures taken after outbreaks		
Vaccination implemented	13	72,22%
Farm blocked	3	16,67%
Change of pasture	1	5,56%
Vaccination recommended (but not done)	1	5,56%

To those veterinarians who saw sporadic cases only (defined as less than five per career so far) we asked specific questions about the place of the occurrence as well as if the case was confirmed by a laboratory or not. Veterinarians seeing lower number of blackleg cases, remembered many specific details about single cases. In total, 12 sporadic cases were described by eight veterinarians. In 50 % of the cases, these occurred on alpine pastures. Twenty-five percent of the cases were located in the valley, the remaining 25 % occurred on pastures, located between the valley and the alpine pasture, where in some regions cattle are kept during May and June before being moved to the final alpine pastures (Table 2).

Measures that were taken after the occurrence of a blackleg case were mainly vaccination of the remaining cattle (72,22 %) (Table 1). None of the veterinarians observed a case in cattle, which had previously been vaccinated.

Vaccination

Sixteen out of 18 veterinarians (88,8 %) regularly performed vaccinations during the last two years (Table 3). The range of applied dosages varied widely from 8 to 1000. Eight veterinarians (44,44 %) used 11–100 vaccine dosages per year, six veterinarians (33,33 %) used 101–300 dosages per year. The main reasons for vaccination were the occurrence of cases in this area in the past (44,44 %), cattle being transferred to summer pastures in risk areas (38,89 %) as well as prophylactic measures in case of known environmental changes such as soil excavations (16,67 %, Table 3).

Most of the interviewed veterinarians (77,78 %), however, reported a decrease in vaccination since 2009, as well as after 2022 (72,22 %), despite them still recommending the prophylactic vaccination to the farmers. Only one participant noted an increase in vaccinations since 2022 due to cases occurring in their practice area during that time (Table 3). The number of animals in alpine pastures remained constant over this time period.³

The majority of participants reported similar arguments against the use of the vaccine, which were mostly mutually dependent. The most frequently mentioned reason against use was the decreasing number of blackleg cases (38,46 %). Furthermore, the need for two injections was mentioned as a reason for denying immunization. Veterinarians reported that it was not always possible to perform a second immunization before moving cattle to the alpine pasture and applying a second booster vaccination on the alpine pasture would have been a huge effort. The occurrence of side effects is another important aspect (23,08 %), especially for farmers participating with their cattle at shows. Local swellings (66,67 %) and abscesses at injection sites (23,81 %) were reported as the most frequent side effects (Table 3). Two participants additionally named reduced general condition of the animal or fever. These findings correspond to the manufacturer's specifications.¹⁰

Discussion

In summary, our results show that most of the officially reported blackleg cases occurred in the areas defined as risk areas. However, some cases also occurred in the areas that were no longer included in the list of risk areas of the Canton of Bern since 2021. This indicates the difficulties of predicting and defining risk areas accurately. Moreover, since the disease is no longer actively monitored, and cases still occur in the Canton of Bern, it is likely that the level of knowledge on previous and future risk areas will decrease. Currently, risk areas in the Canton of Bern are mainly limited to the Bernese Oberland and the northern area of Bernese Jura-Seeland. Other regions (Bern-Mittelland and Emmental-Oberaargau) count less risk areas and have recorded only individual blackleg cases.

Since the disease is no longer reportable, no more official data about the number of cases is available from 2023 onwards. In the Institute of Veterinary Bacteriology at Vetsu-

Occurrence of blackleg and current preventive measures against outbreaks in the Canton of Bern

L. Käser et al.

Table 2: Answers to section of the questionnaire containing questions of blackleg case frequency in the Canton of Bern with a special focus on sporadic cases. Results are displayed in absolute and relative numbers.

Case occurrence ((n), only sporadic cases)	Count	Percent
Alpine pasture	6	50,00 %
Valley	3	25,00 %
Pre-alpine (Voralp)	3	25,00 %
Confirmation ((n), only sporadic cases)		
Yes	8	66,67 %
No	4	33,33 %

Occurrence of blackleg and current preventive measures against outbreaks in the Canton of Bern

L. Käser et al.

in Bern, in 2023, no cases of *C. chauvoei* were isolated. However, in our survey, four practicing veterinarians noted suspected cases of blackleg in 2023, suggesting that the disease is still prevalent in certain areas of the Canton of Bern.

In parallel, our survey revealed a decreasing willingness of farmers to vaccinate cattle against blackleg. This reduction in prophylaxis, as well as reduced disease awareness and

monitoring, together with the biology of *C. chauvoei*, could pose significant risks for reoccurrence of the disease. Spores of *C. chauvoei* are known to be highly resistant to environmental influences and disinfectants and persist over long periods of up to several decades in different environments.^{1,5,17} Thus, recurrent outbreaks in once contaminated environments are typical. Factors such as soil type, flooding events, annual rainfall as well as recent soil excavations have been associated with the occurrence of blackleg.^{15,17}

Table 3: Answers to section of the questionnaire containing questions regarding a multivalent clostridial vaccine usage against blackleg and number of vaccinated animals as well as reasons for and against vaccination and the potential side-effects. Results are displayed in absolute and relative numbers.

Vaccine usage	Count	Percent
Yes	16	88,89%
No	2	11,11%
Number of vaccinated animals (per year)		
None	2	11,11%
1–10	1	5,56%
11–100	8	44,44%
101–300	6	33,33%
>300	1	5,56%
Vaccinations 2009–2022		
Decreasing	14	77,78%
Constant	4	22,22%
Increasing	0	0%
Vaccinations since 2022		
Decreasing	13	72,22%
Constant	4	22,22%
Increasing	1	5,56%
Recommendation of vaccination		
Yes	13	72,22%
No	5	27,78%
Reasons for use of vaccine		
Occurrence of cases	8	44,44%
Summer pastures	7	38,89%
Environmental changes	3	16,67%
Reasons against use of vaccine		
Decreasing number of cases	5	38,46%
Side effects	3	23,08%
Financial aspect	2	15,38%
Delivery Difficulties	1	7,69%
Two injections	1	7,69%
No interest from farmers	1	7,69%
Observed side effects		
Local swelling	14	66,67%
Abscesses	5	23,81%
Decreased general condition	1	4,76%
Fever	1	4,76%

As one of the goals of this study was to understand the preventive measures undertaken in risk areas, majority of the interviewed participants was practicing in risk areas. Therefore, it is of note that the participants were not randomly selected.

Importantly, it is possible that animals become infected while grazing on contaminated pastures, while disease manifests much later in a different location. If animals are not disposed immediately, postmortem opening of the carcass can lead to contamination of new areas. To prevent this, continuous immune prophylaxis in known risk areas is important. Although this study has focused on the Canton of Bern, similar situations could occur in other areas of Switzerland with large cattle populations and alpine pastures.

Veterinarians practicing in once affected areas, in the Canton of Bern and elsewhere in Switzerland, should be aware of the continuing threat of blackleg disease and act accordingly once outbreaks are suspected.

Occurrence of blackleg and current preventive measures against outbreaks in the Canton of Bern

L. Käser et al.

Acknowledgements

We thank the Federal Food Safety and Veterinary Office for providing the data and for their assistance in answering any questions that arose during this study.

We would also like to thank all veterinarians who participated in the survey.

The study was financed by the Institute of Animal Pathology, Vetsuisse-Faculty, University of Bern, Bern, Switzerland. We declare no conflict of interest.

Présence du charbon symptomatique et mesures préventives actuelles contre les foyers dans le canton de Berne

Le charbon symptomatique est une maladie infectieuse aiguë et mortelle causée par *Clostridium chauvoei*. Elle touche principalement les bovins et se caractérise par une myosite nécro-hémorragique et une myocardite graves. Bien que la maladie ait causé des pertes économiques considérables dans le passé, le nombre de bovins malades a diminué au cours des 30 dernières années. Par la suite, le statut du charbon symptomatique a été officiellement modifié en Suisse et la maladie n'est plus à déclarer depuis 2022. Compte tenu de la persistance potentielle des spores de *C. chauvoei* dans l'environnement, le charbon symptomatique pourrait toutefois être encore présente dans certaines zones du canton de Berne, une région agricole importante et étendue de la Suisse.

Dans cette étude, nous avons évalué la distribution spatio-temporelle du charbon symptomatique de 1992 à 2023 et les mesures prophylactiques actuellement appliquées par les vétérinaires dans les zones à risque du canton de Berne. Nos résultats montrent que la plupart des cas se sont produits dans des zones toujours définies comme zones à risque par l'Office vétérinaire cantonal, mais que certains se sont produits dans des zones qui ne sont plus répertoriées comme zones à risque depuis 2020. Parallèlement, les vétérinaires ont signalé des cas suspects au cours des dernières années ainsi qu'une diminution de l'acceptation de la vaccination par les agriculteurs dans les zones à risque.

La combinaison du manque de surveillance du charbon symptomatique, de la méconnaissance de la maladie, de la

Diffusione del carbonchio sintomatico e attuali misure preventive di diffusione nel Canton Berna

Il Carbonchio sintomatico è una malattia infettiva acuta e letale causata da *Clostridium chauvoei*. Colpisce principalmente il bestiame ed è caratterizzata da grave miosite e miocardite necroemorragica. Sebbene in passato la malattia abbia causato ingenti perdite economiche, negli ultimi 30 anni il numero dei bovini colpiti è diminuito. Di conseguenza, lo stato del carbonchio sintomatico è stato ufficialmente modificato in Svizzera e la malattia non è più soggetta a obbligo di notifica dal 2022. Tuttavia, considerando la potenziale persistenza delle spore di *C. chauvoei* nell'ambiente, il carbonchio sintomatico potrebbe essere ancora prevalente in alcune parti del Canton Berna, una regione agricola importante e vasta della Svizzera.

In questo studio, abbiamo valutato sia la distribuzione spatio-temporale del carbonchio sintomatico nel periodo tra il 1992 e il 2023, sia le misure profilattiche attualmente adottate dai veterinari nelle aree a rischio del Canton Berna. I nostri risultati mostrano che la maggior parte dei casi si è verificata in aree ancora definite a rischio dall'Ufficio Veterinario Cantonale, mentre alcuni casi si sono verificati in aree che dal 2020 non sono più elencate come tali. Parallelamente, i veterinari hanno segnalato casi sospetti negli ultimi anni e una ridotta accettazione della vaccinazione da parte degli allevatori nelle zone a rischio.

Purtroppo, la combinazione tra l'assenza di monitoraggio del carbonchio sintomatico, la ridotta consapevolezza della malattia, la diminuzione della disponibilità degli allevatori a vaccinare gli animali e la lunga persistenza ambientale

Occurrence of blackleg and current preventive measures against outbreaks in the Canton of Bern

L. Käser et al.

diminution de la volonté des agriculteurs de vacciner et de la longue persistance des spores de *C. chauvoei* dans l'environnement représente un risque de réapparition de la maladie. Les vétérinaires qui pratiquent dans les zones anciennement touchées, dans le canton de Berne et ailleurs en Suisse, devraient en être conscients et agir en conséquence lorsque des foyers de charbon symptomatique sont suspectés.

Mots clés: charbon symptomatique, canton de Berne, bovins, *Clostridium chauvoei*, zone à risque, acceptation du vaccin

delle spore di *C. chauvoei* rappresentano un rischio per la ricomparsa della malattia. I veterinari che operano nelle aree precedentemente colpite, nel Canton Berna e altrove in Svizzera, dovrebbero esserne consapevoli e agire di conseguenza in caso di sospetti focolai di carbonchio sintomatico.

Parole chiave: Carbonchio sintomatico, Canton Berna, bestiame, *Clostridium chauvoei*, area a rischio, accettazione del vaccino

Literaturnachweis

- ¹ Abreu CC, Blanchard PC, Adaska JM, Moeller RB, Anderson M, Navarro MA, et al.: Pathology of blackleg in cattle in California, 1991–2015. *J Vet Diagn Invest* 2018; 30(6): 894–901.
- ² Abreu CC, Edwards EE, Edwards JF, Gibbons PM, Leal de Araujo J, Rech RR, et al.: Blackleg in cattle: A case report of fetal infection and a literature review. *J Vet Diagn Invest* 2017; 29(5): 612–621.
- ³ AG I: Identitas Plattform Tierstatistik. Bern, CH: <https://tierstatistik.identitas.ch/de/cattle-summering-canton.html> (accessed 15.07.2024).
- ⁴ Bern K: Überblick über die Regionalgerichte. Bern, CH: <https://www.zsg.justice.be.ch/de/start/ueber-uns/regionalgerichte/ueberblick.html> (accessed 01.07.2024).
- ⁵ Junior CAO, Silva ROS, Lobato FCF, Navarro MA, Uzal FA: Gas gangrene in mammals: a review. *J Vet Diagn Invest* 2020; 32(2): 175–183.
- ⁶ Natur AfLu: Landwirtschaft. Amt für Landwirtschaft und Natur, Bern: <https://www.weu.be.ch/de/start/themen/landwirtschaft.html> (accessed 28.06.2024).
- ⁷ Santos B, Ladeira S, Riet-Correa F, Soares M, Marcolongo C, Sallis E, et al.: Clostridial diseases diagnosed in cattle from the South of Rio Grande do Sul, Brazil. A forty-year survey (1978–2018) and a brief review of the literature. *Pesquisa Veterinária Brasileira* 2019; 39: 435–446.
- ⁸ Useh NM, Nok A, Esievo K: Blackleg in ruminants. *Cab Reviews: Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources* 2006: 1.
- ⁹ Uzal FA: Evidence-based medicine concerning efficacy of vaccination against *Clostridium chauvoei* infection in cattle. *The Veterinary clinics of North America Food animal practice* 2012; 28(1): 71–77, viii.
- ¹⁰ Veterinärpharmakologie und -toxikologie If: Bovilis Bravoxin 10 ad us. vet., Injektionssuspension für Rinder und Schafe. Zürich, CH: <https://www.vetpharm.uzh.ch/tak/00000000/00001728.01> (accessed 28.06.2024).
- ¹¹ Veterinärwesen Af: Schutzimpfung gegen Rauschbrand. Bern, CH: <https://www.weu.be.ch/content/dam/weu/dokumente/avet/de/tiere-halten/575x20-VZ-Rauschbrandgebiete-de.pdf> (accessed 28.06.2024).
- ¹² Veterinärwesen BfLu: Strategischer Einsatz von Impfstoffen bei Rindern, Impfleitfaden für Tierärztinnen und Tierärzte. Bern, CH: <https://www.google.ch/url?sa=t&rct=j&q=&esrc=s&source=web&cd=&ved=2ahUKewjWiZT97P2GAxXRzwIHHUQiA1kQFnoECBUQAQ&url=https%3A%2F%2Fwww.blv.admin.ch%2Fdam%2Fblv%2Fde%2Fdokumente%2Ftiere%2Ftierkrankheiten-und-arzneimittel%2Ftierarzneimittel%2Fstar-antibiotika%2Fimpfleitfaden-strategischer-einsatz-impfstoffe-rinder-fuer-tieraerzte.pdf.download.pdf%2Fstrategischer-einsatz-von-impfstoffen-bei-rindern-impfleitfaden-fuer-tierarztinnen-und-tieraerzte-de.pdf&usg=AOvVaw0a9ObbHAcad4bLe0WjedhW&opi=89978449> (accessed 28.06.2024).
- ¹³ Veterinärwesen BfLu: InfoSM Meldesystem. Bern, CH: <https://www.infosm.blv.admin.ch/info> (accessed 01.07.2024).
- ¹⁴ Weissenrieder FX: 25 Jahre Schutzimpfung gegen Rauschbrand im Kanton Bern und deren Erfolg. – 1895 bis 1920. *Schweizer Archiv für Tierheilkunde* 1921: 12: 524–547.
- ¹⁵ Wolf R, Hiesel J, Kuchling S, Deutz A, Kastelic J, Barkema HW, et al.: Spatial-temporal cluster analysis of fatal *Clostridium chauvoei* cases among cattle in Styria, Austria between 1986 and 2013. *Preventive Veterinary Medicine* 2017: 138: 134–138.
- ¹⁶ Wyss R: Sömmerungsvorschriften 2024 für den Kanton Bern. Amt für Veterinärwesen, Bern: <https://www.weu.be.ch/content/dam/weu/dokumente/avet/de/tiere-halten/434x10-S%C3%B6mmerungsvorschriften-de.pdf> (accessed 28.06.2024).
- ¹⁷ Ziech RE, Gressler LT, Frey J, Vargas ACd: Blackleg in cattle: current understanding and future research needs. *Ciência Rural* 2018: 48.

Korrespondenzadresse

Dr. Iva Cvitas, DVM, PhD, Dipl. ECVP
Länggassstrasse 122
3012 Bern
Telefon: +41 31 684 24 08
E-Mail: iva.cvitas@unibe.ch